

Mestsoort	SG	DS%	OS%	Gehalte in kg/ton								
				N-tot	N-min	N-org	P2O5	K2O	MgO	Na2O	S	
Vloeibare mest												
Rundveedrijfmest	1.005	8.5	6.4	4.1	2.0	2.1	1.5	5.8	1.2	0.7	0.7	
Rosé kalveren	--	9.4	7.1	5.6	3.0	2.6	2.6	5.0	1.6	1.2		
Witvlees kalveren	1.000	2.2	1.7	2.6	2.1	0.5	1.1	4.5	1.7	1.6	0.2	
Vleesvarkensdrijfmest												
Zeugendrijfmest	1.000	6.7	2.5	5.0	3.3	1.7	3.5	4.9	1.4	0.9	0.4	
Kippendrijfmest	1.020	14.5	9.3	10.2	5.8	4.4	7.8	6.4	2.2	0.9	0.9	
Rundveegier	1.030	2.5	1.0	4.0	3.8	0.2	0.2	8.0	0.2	1.0	0.8	
Varkensgier	1.010	2.0	0.5	6.5	6.1	0.4	0.9	4.5	0.2	1.0	0.7	
Zeugengier	1.020	1.0	1.0	2.0	1.9	0.1	0.9	2.5	0.2	0.2	0.2	
Mestsoort	SG	DS%	OS%	N-tot	N-min	N-org	P2O5	K2O	MgO	Na2O	S	
Vaste mest												
Rundvee	0.900	19.4	15.2	5.3	0.9	4.4	2.8	6.1	2.2	1.0	0.2	
Varkens	0.800	26.0	15.3	7.9	2.6	5.3	7.9	8.5	2.5	0.9	0.6	
Leghennen, mestband	0.605	57.3	41.6	25.6	2.5	23.1	19.6	15.5	5.5	1.7	2.5	
Leghennen, mestband+ nadroog	--	81.0	42.7	34.1	3.9	30.2	27.8	20.1	5.9	2.3		
Kippen strooiselmest	0.600	71.3	35.9	28.0	3.6	24.4	25.6	20.8	7.5	3.4	3.6	
Vleeskuikens+parelhoen	0.600	62.6	41.9	32.1	8.0	24.1	16.8	20.5	7.1	3.0	3.6	
Kalkoenen	0.530	52.0	42.7	23.3	6.0	17.3	19.7	13.4	5.8	6.7	3.8	
Eenden	0.900	27.5	23.7	8.9	1.6	7.3	7.3	8.4	3.4	1.3	2.0	
Konijnen	0.600	40.8	33.2	9.4	2.3	7.1	6.7	10.7	5.2	2.0	3.0	
Paarden	0.700	28.7	16.0	4.6	0.5	4.1	2.7	8.1	1.8	1.6	2.0	
Schapen	0.700	29.0	20.6	8.6	2.0	6.6	4.2	16.0	2.8	2.3	1.9	
Nertsen, Vossen	--	45.2	29.3	28.3	16.1	12.2	26.9	5.4	3.5	8.1	5.9	
Geiten	0.900	29.1	17.4	9.9	2.4	7.5	5.3	12.8	4.0	1.9	2.0	
Champost	0.550	30.0	17.7	5.2	0.3	4.9	3.2	8.7	2.4	0.9	5.5	
GFT-compost	0.800	61.0	18.0	6.9	0.7	6.2	3.5	6.8	3.3	1.9	3.8	

Bron: Nutrinorm